



ISSN 2395-650X

International Journal of
Life Sciences Biotechnology Pharma Sciences

IJLBPS

www.ijlbps.org

E-mail: editorijlbps@gmail.com editor@ijlbps.org

New gastroretentive medication delivery method using in situ gel **CH.Raju**

Due to their rapid elimination from the stomach and duodenum, liquid dose forms are more likely to have poor bioavailability than other oral dosage forms. A liquid in-situ gelling device might be an effective way to increase the production of a sustained release formulation of an oral liquid formulation. Gels may be formed in response to variations in temperature, pH, ionic strength, and ultraviolet light, all of which influence the rate and duration of drug release. Advantages of in situ gel forming polymeric formulations include increased bioavailability of medicine and patient compliance via decreased dose frequency, as well as sustained and extended activity in contrast to traditional drug delivery methods. Keywords: In situ gel, Sustained drug delivery, Ionic crosses linking.

Introduction

There has been a lot of research on in situ gel forming systems as potential vehicles for long-term drug delivery. The benefits of in situ forming polymeric delivery methods, such as simple administration, decreased dosing frequency, increased patient compliance, and comfort, have inspired this interest.¹ Changes in pH, temperature, and solvent exchange² are all triggers that may lead to in situ gel formation. This allows for the development of In-Situ Gelling Systems for administration through various routes (oral, nasal, ophthalmic, etc.). Gellan gum, alginic acid, xyloglucan, pectin, chitosan, poly (DL lactic acid), poly (DL- lactide-co-glycolide), and polycaprolactone are only some of the natural and synthetic polymers employed in the formulation creation of in situ forming drug delivery systems.

Compared to the standard liquid dose form, the bioavailability of the medicine is improved by the gastroretentive in situ gelling mechanism. Since the gel formed by the in situ gelling system is less dense than the gastric fluids, it is able to float on top of the stomach contents or adhere to the gastric mucosa thanks to the polymer's bioadhesive properties, resulting in the prolonged delivery of the drug. This article aims to explore the aspects to be addressed in the formulation of an in-situ drug delivery system, with a focus on stomach-specific in situ gelling systems. Evaluation and characterisation of in-situ polymeric formulations, as well as the processes of gel formation from sol forms of various smart polymers, are also covered. The upsides of using a GRDDS (gastric-retention drug-delivery system).

The GRDDS framework is applicable to any drug or group of drugs.

medications that are absorbed in the stomach, such as ferrous salts, and medications that are intended for local action in the stomach, such as antacids, benefit from the GRDDS. Medications' effectiveness may be improved with the help of the continuous release. Poor absorption is to be anticipated when intestinal movement is rapid and transit time is short, as may occur in some types of diarrhea. In these cases, the medicine may be more effective when retained in the stomach by the process of gastroretention.

Drugs with tiny intestine absorption windows can be delivered using GRDDS, which is a benefit. Medicines absorbed primarily via the stomach are not the only substances affected by the GRDDS. This is because it

has been discovered that they are just as effective as orally ingested drugs. chlorpheniramine maleate, for instance. Medications that might work well in a gastroretentive dosage form. Specific gastrointestinal (GI) absorption window, such as for riboflavin and levodopa. Absorption occurs mostly in the stomach and upper GI tract; examples include calcium supplements, chlorthalidone, and cinnarizine. Locally acting drugs in the stomach, such as antacids and misoprostol. Drugs include ranitidine HCl and metronidazole that break down in the gut. Medications like amoxicillin trihydrate that kill off healthy colonic flora. Methods that take use of in-situ gelling systems. In situ gel formation may be triggered by a variety of processes, including chemical reactions (such as enzymatic, chemical, and photo-initiated polymerization), physiological stimuli (such as temperature and pH), and physical changes in biomaterials (such as diffusion of solvent and swelling). Physically based in situ formation. Inflation and Dispersion Gel is formed when a polymer absorbs water and expands. Myverol 18-99 (glycerol mono-oleate), a biodegradable lipid molecule, produces an in situ gel under such conditions.⁷ Diffusion of the solvent from the polymer solution into the surrounding tissue causes

the polymer matrix to precipitate or solidify when using a polymer such as N-methyl pyrrolidone (NMP).⁸ Gelation triggered on-site by chemical agents. Cross-linking of ions. Phase change occurs in ion-sensitive polysaccharides such as carrageenan, Gellan gum (Gelrite®), pectin, and sodium alginate. As a result of the presence of ions like K^+ , Ca^{2+} , Mg^{2+} , and Na^+ .⁹ Because of its interaction with the guluronic acid block in alginate chains, alginic acid gels when divalent/polyvalent cations like Ca^{2+} are present.¹⁰ Cross-linking by enzymes. The rate of gel formation can be easily controlled by using specific natural enzymes that function well under physiological conditions without the use of potentially harmful chemicals like monomers and initiators, allowing the mixtures to be injected before gel formation in situ.¹¹

Photo-polymerisation

The application of electromagnetic radiation can be used to form gel designed readily to be degraded by chemical or enzymatic processes or can be designed for long term persistence in vivo from a solution of monomers such as acrylate or other polymerizable functional groups and initiator such as 2,2 dimethoxy-2-phenyl acetophenone, camphorquinone, and ethyl eosin injected into a tissues site.¹² The utilization of UV and visible light with extended wavelengths is common. As a tissue-contacting material and controlled-release carrier, Sawhney¹³ reports the use of a photopolymerizable, biodegradable hydrogel. Formation of gels on the spot in response to physiological cues. In situ gelation is temperature dependent. Before administration, these aqueous solutions are liquid, but they solidify when heated to body temperature. These hydrogels are liquid at room temperature (about 20°C - 25°C) but gel when exposed to bodily fluids (around 35°C - 37°C). This method takes use of a phase transition that is temperature-dependent. The solubility of certain polymers suddenly shifts when the surrounding temperature rises (the polymer's LCST).^{14,15} In contrast to polymer-polymer and



water-water interactions, hydrogen bonding between the polymer and water becomes unfavorable at the LCST, causing a sudden transition as the solvated macromolecule rapidly dehydrates and transitions to a more hydrophobic structure.^{16,17} On the other hand, certain amphiphilic polymers, which self-assemble in solution, exhibit micelle packing and gel formation at elevated temperatures due to polymer-polymer interactions.¹⁸ Among environment-sensitive polymer systems, temperature-sensitive hydrogels get the greatest attention in the field of drug delivery.¹⁹ Pluronics (poly (ethylene oxide)-poly (propylene oxide)-poly (ethylene oxide) (PEO-PPOPEO) Triblock) polymers are one example.²⁰ Networks of polymers consisting of poly(acrylic acid), poly(acrylamide), or poly(acrylamide-co-butyl methacrylate). At room temperature, polymer solutions flow freely but gel when near the human body.²² Hydrogels with a positive temperature coefficient shrink when cooled below their upper critical solution temperature (UCST). There is a positive temperature dependency of swelling in polymer networks made of poly (acrylic acid) (PAA) and polyacrylamide (PAAm) or poly (acrylamide-co-butyl methacrylate).²¹ Gelling that depends on pH. In addition, a shift in pH may trigger the development of in situ gel. Poly (methacrylic acid) (PMA) and poly (ethylene glycol) (PEG) mixtures²⁵ and PAA (Carbopol®, carbomer) or its derivatives²³ demonstrate a transition from sol to gel with a change in pH. Polyelectrolytes are polymers that include several ionizable groups. If the polymer has weakly acidic (anionic) groups, the swelling of the hydrogel will increase as the external pH rises, whereas it will decrease if the polymer contains weakly basic (cationic) groups. Gelling system polymers for use in the mouth. Oral medication delivery systems are formed in situ using natural polymers such pectin, xyloglucan, sodium alginate, and gellan gum.

Pectin

Almost all plant cell walls include pectins, which are anionic polysaccharides. Pectin has -(1-4)-D-galacturonic acid residues as its backbone. (See




Figure 1). When divalent ions such free calcium ions are present, the galacturonic acid chains are crosslinked in an egg-box fashion, causing the aqueous solution to solidify.²⁶ When taken orally, pectin undergoes a phase change to the gel state due to the presence of H^+ ion.²⁷ For the purpose of inducing pectin gelation, the formulation may incorporate calcium ions in the complexed form. The degree of esterification (DE) of commercially available pectins ranges from low (LM) to high (HM). When divalent ions, such Ca^{2+} , are present, LM pectins gel, but when the pH drops below roughly 3.3, they may also gel without the presence of Ca^{2+} . In situ pectin gel was created by Kubo and colleagues, Pectin aqueous solutions (1.0 and 1.5%, w/v) were tested as carriers for the prolonged release of the expectorant medication Ambroxol hydrochloride because of their ability to form in situ gels after oral administration. Pectin gelation was triggered by the release of calcium ions from their complexes in the stomach's acidic environment. Ambroxol was shown to be released from the gels at a regulated pace by diffusion in in vitro trials spanning 6 hours. Gels containing an identical dose of Ambroxol formed in situ in the stomachs of rats achieved a bioavailability of approximately 64% of that of a commercially available formulation, with appreciably lower peak plasma levels and a sustained release of drug for at least 6 h. Rheological and drug release features of the formulations have been studied to determine the impact of additional sorbitol (17%, w/v).





Fig. 1: Structure of pectin
Xyloglucan

Xyloglucan is a polysaccharide derived from tamarind seeds and is composed of a (1-4)- β -D-glucan backbone chain, which has (1-6)- α -D xylose branches that are partially substituted by (1-2)- β -D-galactoxylose³¹ Xyloglucan is composed of heptasaccharide, octasaccharide and nonasaccharide oligomers, which differ in the number of galactose side chains.³² (Fig. 2)

Fig. 2: Backbone structure of xyloglucan. Glc,xyl,Gal indicate β -D xylopyranosyl and β -D-galactopyranosyl residues respectively



Although xyloglucan itself does not gel, dilute solutions of xyloglucan which has been partially degraded by galactosidase exhibit a thermally reversible sol-gel transition on heating. Xyloglucan gels have potentially been used for oral, intraperitoneal, ocular and rectal drug delivery.^{33, 34, 35, 36} Its potential application in oral delivery exploits the

proposed slow gelation time (several minutes) that would allow in situ gelation in the stomach following the oral administration of chilled xyloglucan solution. Itoh K examined the gelation and release characteristics of mixtures of xyloglucan, which has thermally reversible gelation characteristics, and pectin, the gelation of which is ion responsive, with the aim of formulating an in situ gelling vehicle suitable for oral sustained drug delivery.³⁷

Gellan gum

Gellan gum (commercially available as Gelrite™ or Kelcogel™) is an anionic deacetylated exocellular polysaccharide secreted by *Pseudomonas elodea* with a tetrasaccharide repeating unit of one α -L-rhamnose, one β -D-glucuronic acid and two β -D-glucuronic acid residues³⁸ Chemical structure of the polysaccharide has a tetrasaccharide repeat unit consisting of two glucose (Glc) residues, one glucuronic acid (GlcA) residue, and one rhamnose (Rha) residue. These are linked together to give a tetrasaccharide repeat unit (Fig. 3)³⁹

Fig. 3: Structure of gellan gum

Toxicological study of Gellan gum has been performed in rats. Male and female Sprague-Dawley rats (20/sex/group) were fed dietary levels of Gellan Gum ranging from 0-6% for 13 weeks. Although the animals on this study experienced symptoms of a sialodacryoadenitis viral infection, all animals survived treatment and there were no adverse effects associated with the feeding of Gellan Gum.⁴⁰ Gellan gum produces temperature dependent or cations induced in situ gelling⁴¹. An increased bioavailability with

sustained drug release profile of theophylline in rats and rabbits was observed from gellan formulations as compared to the commercial sustained release liquid dosage form.⁴⁰ Effect of Mg^{2+} and Ca^{2+} on gel hardness has been shown in following graph. (Fig. 4)





Fig. 4: Effect of divalent cations on gel Hardness of 1% Gelrite gel

Rajinikanth prepared stomach specific in situ gelling system of Clarithromycin to eradicate *H. pylori* infection using gellan gum as a polymer. Gellan based formulation was prepared by dissolving varying concentrations of gel



lan in deionized water to which varying concentrations of drug and sucralfate were dispersed well. The formulation parameters like concentrations of gellan gum and sucralfate influenced the rate and extent of in vitro drug release significantly from FIGC. The addition of sucralfate to the formulation significantly suppressed the degradation of clarithromycin at low pH.⁴²

Sodium Alginate

Sodium alginate is a salt of Alginic acid - a linear block copolymer polysaccharide consisting of β -D-mannuronic acid and α -L-glucuronic acid residues joined by 1,4-glycosidic linkages.⁴³ (Fig. 5). Aqueous solutions of alginates form

firm gels on addition of di- and trivalent metal ions. Nakamura et al. investigated the thermal properties of water insoluble alginate films containing di and trivalent cations. The results indicated that the alginates form compact structures when the ionic radii of the cation are lower. Changes in the film structure during ionic exchange were studied on the basis of its glass transition temperature (T_g) and heat capacity using differential scanning calorimetry (DSC).⁴⁴ Sodium alginate has been employed in the preparation of gels for the delivery of biomolecules such as drugs, peptides and proteins.⁴⁵ Rohit et al., developed a gastroretentive in situ gelling liquid formulation for controlled delivery of ranitidine using sodium alginate (low, medium and high viscosity grades), calcium carbonate (source of cations) and ranitidine. Prepared formulations were evaluated for viscosity, buoyancy lag time and buoyancy duration, drug content and in vitro drug release. Formulation variables such as concentration of sodium alginate, calcium carbonate and drug significantly affected the formulation viscosity, floating behavior and in vitro drug release. Analysis of the release pattern showed that the drug release from in situ gel followed a diffusion mechanism⁴⁶. It exhibits favorable biological properties such as biodegradability and nontoxicity⁴⁷ and mucoadhesive properties. Evaluation of in situ gelling system ClarityThe clarity of formulated solutions can be determined by visual inspection under black and white background.

Viscosity

This is an important parameter for the in situ gels, to be evaluated.

Fig. 5: Structure of Alginic acid

The viscosity and rheological properties of the polymeric formulations, either in solution or in gel made with artificial tissue fluid (depending upon the route of administrations) were determined with different viscometer

The viscosity of these formulations should be such that it should be patient compliant. Sol-Gel transition temperature and gelling time For in situ gel forming systems, the sol-gel transition



temperature and pH should be determined. Gelling time is the time required for first detection of gelation of in situ gelling system.³⁶ Thermosensitive in situ gel should be checked for in situ gelling at body temperature.

Gel-Strength A specified amount of gel is prepared in a beaker, from the sol form. This gel containing beaker is raised at a certain rate, so pushing a probe of rheometer slowly through the gel. The changes in the load on the probe can be measured as a function of depth of immersion of the probe below the gel surface.³⁶ Fourier transform infra-red spectroscopy and Thermal analysis

Fourier transform infra-red spectroscopy is performed to study compatibility of ingredients. Differential scanning calorimetry is used to observe if there are any changes in thermograms as compared with the pure ingredients used thus indicating the interactions.⁵⁰

In-vitro drug release studies

The drug release studies are carried out by using the plastic dialysis cell. The cell is made up of two half cells, donor compartment and a receptor compartment. Both half cells are separated with the help of cellulose membrane. The sol form of the formulation is placed in the donor compartment. The assembled cell is then shaken horizontally in an incubator. The total volume of the receptor solution can be removed at intervals and replaced with the fresh media. This receptor solution is analyzed for the drug release using analytical technique.

Conclusion

In situ gelling system becomes helpful as an alternative of oral solid dosage form with an advantage of liquid dosage form. Sustained release formulation can be prepared in liquid form using in situ gelling approach. In situ gelling system not only helpful for sustained drug delivery, but also become convenient for pediatric and geriatric patient. Exploitation of polymeric in-situ gels for controlled release of various drugs provides a number of advantages over conventional dosage forms. Good biocompatibility characteristics also make the in situ gel dosage forms very reliable.

References

- Peppas N. and Langer R. (1994). New challenges in biomaterials. *Science*, **263**:1715-1720.
- Sarasija S. and Shyamala B. (2005). Nasal Drug Delivery: An Overview. *Indian J. Pharm. Sci.*, **67**: 19-25.
- Wataru K., Yasuhiro K., Miyazaki S. and Attwood D. (2004). In situ gelling pectin formulations for oral sustained delivery of paracetamol. *Drug Develop. Ind. Pharm.*, **30**:593-602.
- Whitehead L., Fell J. and Collett J. (1996). Development of a Gastroretentive Dosage Form. *Eur. J. Pharm.Sci.*, **4**:182-187.
- Deshpande A., Shah N., Rhodes C. and Malick W. (1997). Development of a novel controlled release system for gastric retention. *Pharm Res*, **14**: 815-824.
- Esposito E. and Carratto V. (1996). Comparative analysis of tetracycline containing dental gels; poloxomers and mono-glycerides based formulation. *Int. J. Pharm*, **142**:9-23.
- Geraghty P. and Attwood D. (1997). An investigation of parameters influencing the bioadhesive properties of myverol 18-99/ water gels. *Biomaterials*, **18**:63-70.
- Motto F. and Gailloud P. (2000). In-vitro assessment of new embolic liquids prepared from preformed polymers and water miscible solvents aneurysm treatment. *Biomaterials*, **21**:803-811.
- Bhardwaj T.R., Kanwar M., Lal R. and Gupta A. (2000). Natural gums and modified natural gums as sustained release carriers. *Drug Devel. Ind. Pharm.*, **26**:1025-1038.
- Guo J., Skinner G., Harcum W. and Barnum P. (1998). Pharmaceutical applications of naturally occurring water soluble polymers. *Pharm Sci. & Technol. Today*, **1**:254-261.
- Podual K. and Peppas N.A. (2000). Dynamic behavior of glucose oxidase-containing microparticles of poly(ethylene)- grafted cationic hydrogels in an environment of changing pH. *Biomaterials*, **21**:1439-1450.
- Burkoth A. and Anseth K. (2000). A review of photocrosslinked polyanhydrides: In situ forming degradable networks. *Biomaterials*, **21**:2395- 2404.
- Sawhney A. and Pathak C. (1995). Photopolymerizable biodegradable hydrogels as tissue contacting materials and controlled release carriers. US Patent 5410016.
- Taylor L. and Cerankowski L. (1975), Preparation of films exhibiting a balanced temperature dependence to permeation by aqueous solutions—a study of lower consolute behavior. *J. Polym. Sci. Polym. Chem. Ed.*, **13**: 2551–2570.



- Heskins M. and Guillet J. (1968). Solution properties of poly (N-isopropylacrylamide). *J. Macromol. Sci. Chem.*, **2**: 1441–1455.
- Schild H. (1992). Poly (N-isopropylacrylamide): experiment, theory and application. *Prog. Polym. Sci.*, **17**: 163-249.
- Hoffman A. (1997). *Intelligent polymers*, In: K. Park (Ed.), *Controlled Drug Delivery—Challenges and Strategies*. American Chemical Society, Washington, 485–498.
- Mortensen K. and Pederson J. (1993). Structural study on the micelle formation of poly (ethylene oxide)-poly (propylene oxide)-poly (ethylene oxide) triblock copolymer in aqueous solution. *Macromolecules*, **26**: 805–812.
- Jones M. (2002). In-situ forming biomaterials. *Surg. Clin. N. Am.*, **14**:29-38.
- Peppas N., Bures P., Leobandung W. and Ichikawa H. (2000). Hydrogels in pharmaceutical formulations. *Eur. J. Pharm. Biopharm.*, **50**:27-46.
- Qiu Y. and Park K. (2001). Environment-sensitive hydrogels for drug delivery. *Adv. Drug Deliv. Rev.*, **53**:321-339.
- Bromberg L. and Ron E. (1998). Temperature-responsive gels and thermogelling polymer matrices for protein and peptide delivery. *Adv. Drug Deliv. Rev.*, **31**:197-221.
- Soppimath K., Aminabhavi T., Dave A, Kumbar S. and Rudzinski W. (2002). Stimulus-responsive smart hydrogels as novel drug delivery systems. *Drug Dev. Ind. Pharm.*, **28**: 957- 974.
- Aikawa K., Mitsutake A., Uda H., Tanaka S., Shimamura H. and Aramaki Y.(1998). Drug release from pH-response polyvinylacetal diethyl aminoacetate hydrogel, and application to nasal delivery. *Int. J. Pharm.*, **168**:181-189.
- Alexandridis P. and Lindman B. (2000). *Amphiphilic block polymers*. Amsterdam:Elsevier.
- Dumitriu S., Vidal P.F. and Chornet E. (1996). Hydrogels based on polysaccharides. In: Dumitriu S., Editor. *Polysaccharides in medical applications*. New York: Marcel Dekker Inc, 125–242.
- Ni Y. and Kenneth M.Y. (2004). In-situ gel formation of pectin. United States Patent 6777000.
- Thakur B., Singh R. and Handa A. (1997). Chemistry and uses of pectin—A review. *Crit. Rev. Food Sci.Nutr.*, **37**: 47–73.
- Gilsenan P., Richardson R. and Morris E. (2000). Thermally reversible acid induced gelation of low-methoxy pectin. *Carbohydr. Polym.*, **41**: 339–349.
- Kubo W and Miyazaki S (2004). Oral sustained delivery of ambroxol from in situ-gelling pectin formulations. *Int J Pharm.*, 271:233-40.
- Miyazaki S. and Kawasaki N. (2001). Comparison of in situ gelling formulations for the oral delivery of cimetidine. *Int. J. Pharm.*, **220**:161–168.
- York W. Van H., Darvill A. and Albersheim P. (1990). Structural analysis of xyloglucan oligosaccharides by ¹H-n.m.r. spectroscopy and fast atom-bombardment mass spectrometry. *Carbohydr. Res.*, **200**: 9–31.
- Kawasaki N., Ohkura R., Miyazaki S., Uno Y., Sugimoto S. and Attwood D. (1999): Thermally reversible xyloglucan gels as vehicles for oral drug delivery. *Int. J. Pharm.*, **181**:227–234.
- Suisha F., Kawasaki N., Miyazaki S., Shirakawa M., Yamotoya K. and Sasaki M. (1998). Xyloglucan gels as sustained release vehicles for intraperitoneal administration of mitomycin C. *Int. J. Pharm.*, **172**:27–32.
- Miyazaki S., Suzuki S., Kawasaki N., Endo K., Takahashi A. and Attwood D. (2001). In situ gelling xyloglucan formulations for sustained release ocular delivery of pilocarpine hydrochloride. *Int. J. Pharm.*, **229**:29–36.
- Miyazaki S., Suisha F. and Kawasaki N. (1998). Thermally reversible xyloglucan gels as vehicles for rectal drug delivery. *J. Control Rel.*, **56**:75–83.
- Itoh K. and Yahaba M. (2008). In situ gelling xyloglucan/pectin formulations for oral sustained drug delivery. *Int. J. Pharm.*, **356**: 95-101.
- Miyazaki S., Hirotatsu A., Kawasaki N., Wataru K. and Attwood D. (1999). In situ gelling gellan formulations as vehicles for oral drug delivery. *J. Control Rel.*, **60**:287–295.



- Jansson P. and Lindberg B. (1983). Structural studies of gellan gum, an extracellular polysaccharide elaborated by *Pseudomonas elodea*. *Carbohydr. Res.*, **124**: 135-139.
- Batham P., Rainey S., Bier C., Losos G., Osborne B. and Procter B. (1983). A 13-week toxicity study of a polysaccharide gum (K9A50) during dietary administration to the albino rat. Bio -Research Laboratories Ltd., Montreal, Canada.
- Crescenzi V., Dentini M. and Coviello T. (1990). Solutions and gelling properties of microbial polysaccharides of industrial interest. Novel biodegradable microbial polymers: The case of gellan. Dordrecht: Kluwer Academic Publishers, In Dawes E.A., editor, 227–84.
- Rajinikanth P. and Mishra B. (2008). Floating in situ gelling system for stomach site-specific delivery of clarithromycin to eradicate *H. pylori*. *Journal of Controlled Release*, **125**:33-41.
- Sechoy O., Tissie G., Sebastian C., Maurin F., Driot J.Y. and Trinquand C. (2000). A new long acting ophthalmic formulation of carteolol containing Alginic acid. *Int. J. Pharm.*, **207**:109–116.
- Nakamura K., Nishimura Y., Hatakeyama H. and Hatakeyama T. (1995). Thermal properties of water insoluble alginate films containing di- and trivalent cations. *Thermochim. Acta*, **267**:343-353.
- Wong W., Chan L., Kho S. and Heng P.W. (2002). Design of controlled-release solid dosage forms of alginate and chitosan using microwave. *J. Controlled Release*, **84**: 99-114.
- Rohith G., Sridhar B.K. and Srinatha A. (2009). Floating drug delivery of a locally acting H₂-antagonist: an approach using an in situ gelling liquid formulation. *Acta Pharm.*, **59**:345-54.
- Al-Shamklani A., Bhakoo M., Tuboku M.A. and Duncan R. (1991). Evaluation of the biological properties of alginates and gellan and xanthan gum. *Proc. Int. Symp. Control Release Bioact. Mater.*, **18**:213–217.
- Smart J., Kellaway I. and Worthington H. (1984). An in vivo investigation of mucosa adhesive materials for use in controlled drug delivery. *J. Pharm Pharmacol.*, **36**:259–299.
- Fuonfuchat A., Jamieson A., Blackwell J. and Gerken T.A.(1996). Rheological studies of interaction of mucins with alginates and polyacrylate. *Carbohydr Res.*, **284**:85–99.
- Kashyap N., Viswanad B., Sharma G., Bhardwaj V., Ramarao P. and Kumar M.N. (2007). Design and evaluation of biodegradable, biosensitive in situ gelling systems for pulsatile delivery of insulin. *Biomaterials*, **28**:2051-2060.