



ISSN 2395-650X

**International Journal** of  
Life Sciences Biotechnology Pharma Sciences

IJLBPS

[www.ijlbps.org](http://www.ijlbps.org)

E-mail: [editorijlbps@gmail.com](mailto:editorijlbps@gmail.com) [editor@ijlbps.org](mailto:editor@ijlbps.org)



## Dye industry effluent treatment using free and immobilized fungus

A.Arivanbabu

---

### Abstract

Paper and pulp mills, textile and dye factories, distilleries, and the leather industry are just few of the businesses that produce brightly colored effluents. The lignin-degrading enzymes laccase, manganese-peroxidase, and lignin peroxidases produced by terrestrial white-rot basidiomycetous fungus may be used to neutralize dyes and other xenobiotics in industrial effluents. Thirteen unique fungal species were identified from the wastewater for this investigation. *Aspergillus* was found to be the most common fungal genus in the waste water. In comparison to the control group, all variables except dissolved oxygen exhibited a reduction. Both the BOD and the COD levels dropped dramatically. *Aspergillus* that has been immobilized was more effective in cleaning up pollution than free-living *Aspergillus* cells.

Key-Words: *Aspergillus*, BOD, COD

---

### Introduction

It has been predicted that by 2010 the United States would produce 253 million tons of municipal solid garbage annually (USEPA, 2007). With a growing rate of roughly 3% per year (PCD, 2005)[1], solid waste production in Thailand's metropolitan regions exceeds 14.3 million tons per year, or 39,200 tons per day. About 65% of garbage is disposed of in open dumps while the remaining 35% is handled via sanitary landfill. Basidiomycete fungi are well-known for producing ligninolytic enzymes. White rot fungus's fermented culture broth is rich in lignin modifying enzymes (LMEs) [2,3], whereas bacteria, yeast, and other fungi seldom make LMEs. The efficacy of treating Olive Oil mill wastewaters (OOMW) with four different white rot fungi (WRF) strains was evaluated in relation to their cultivation mode, specifically as free mycelium, mycelium immobilized in alginate beads, and solid state cultivation on Petri dishes[4].

---

Department of Microbiology, Department of Chemistry, Department of Biochemistry. Shrimati Indira

Gandhi College, Trichy, (TN) - India

---



## Material and methods

### *Isolation of fungi*

Ten ml of the effluent sample was taken in a 250ml conical flask containing 90ml sterile distilled water. The flask was shaken on an elastic shaker to get a homogenous suspension and transferring serially 10ml of the effluent suspension to 90ml of sterile distilled water made different dilutions viz., 10<sup>-1</sup>, 10<sup>-2</sup> and 10<sup>-3</sup>.

3. 0.1 ml of 10<sup>-3</sup> dilution was plated in petridishes containing Potato Dextrose Agar medium (PDA). **Immobilization of fungi in Alginate beads** (Fig.2) Commercial grade sodium alginate was used as the immobilizing agent for bead preparation. Exponentially growing *Aspergillus flavus* (Fig.1) were harvested by centrifugation (5000 x g for 10 min) and resuspended in 50ml of sterile water. To this 50ml of 4% alginate solution was mixed thoroughly to get a final alginate concentration of 2%. The alginate-bacterial mixture was then added drop wise into CaCl<sub>2</sub> (0.1 M) solution. The beads were kept in the same solution for 30 mins at 4°C for hardening.

### **Experimental condition** (Fig.3)

For the present study the following treatments were employed

- Effluent sample without inoculation-control I
- Effluent sample inoculated with *Aspergillus flavus*-treatment I
- Effluent sample inoculated with immobilized *A. flavus*- treatment II

The experiment was conducted in patch cultures in duplicates for a total period of 15 days in 250ml Erlenmeyer flasks. Effluent samples (control and treated) were periodically (every 5<sup>th</sup> days) analyzed for various physico-chemical parameters and recorded.

### **Physico-chemical analysis of effluent** (APHA, 1995)

#### *Determination of pH*

pH was recorded at the collection site with pH indicator paper. At laboratory the pH was checked again with pH meter (Elico-India).

#### *Estimation of alkalinity (Carbonate and Bicarbonate)*

To 10ml of filtered sample, 2 drops of phenolphthalein indicator was added and titrated

against 0.1N HCl. This sample was used for both phenolphthalein and total alkalinity.

#### *Estimation of Free Carbon dioxide*

To 10ml of filtered sample a few drops phenolphthalein indicator was added and titrated against 0.05 N NaOH. The end point was the appearance of pink color.

#### *Estimation of Dissolved oxygen*

Standard flasks (15 ml) were filled with filtered samples without air bubbles. Then the samples were Winklerised by adding 0.2 ml of manganous sulphate followed by 0.2 ml of alkaline iodide. The precipitate was dissolved by the addition of 0.2 ml of sulphuric acid. The clear yellow color solution was titrated against sodium thiosulphate and starch was used as an indicator. The end point was the disappearance of blue color.

#### *Estimation of Nitrate*

A series of 50ml standard solutions of the nitrate from the standard nitrate solution were prepared to obtain a range of conc. (10 to 100µg). 2ml aliquots of the standard solutions were pipetted into different dry 100ml beakers provided with glass rods. To each beaker 1ml brucine sulphanilic acid reagent and 10ml sulphuric acid solution was added. Stirred gently for about 5 minutes. The beakers were covered with the watch glasses and kept in dark for 10 minutes during which, a yellow colour develops. 10ml distilled water was added to each of the beakers and incubated in the dark for 30 min. The absorbance of each solution was then measured at 410nm. Plotting the absorbance values against the nitrate concentrations of the standard solutions draws a graph. The amount of nitrate content of the effluent was determined using the standard graph.

#### **Estimation of Nitrite**

To 10ml of filtered sample, 0.2ml of sulphanilamide solution was added. After 10 minutes 0.2ml of NED was added. The pink colour appeared was measured at 543nm in spectronic 20 against reagent blank. The amount of Nitrite (mg/l) was calculated from the standard graph prepared by using sodium nitrite.

#### *Estimation of Ammonia*

To 10ml of filtered sample 0.4ml phenol reagent was added and stirred, then 0.4ml of nitroprusside reagent was added and mixed. Finally, 1ml of the oxidizing reagent was added and mixed thoroughly and then incubated at room temperature for an hour. Absorbency was measured at 630nm in spectronic 20. The amount of ammonia was calculated by using a standard graph prepared by using ammonium chloride.

#### *Estimation of total phosphorus*

To 10ml of filtered sample and 2 ml of 5% potassium persulphate followed by 2ml of mixed reagent was added and diluted to 15ml with distilled water. The reading was taken after 10 minutes in spectronic 20 against the reagent blank at 882 nm. The amount of inorganic phosphate was calculated using a standard graph prepared by using  $K_2HPO_4$ .

#### *Estimation of inorganic phosphate*

To 10ml of filtered sample, 2ml of mixed reagent was added and diluted to 15ml with distilled water. The reading was taken after 10 minutes in spectronic 20 against the reagent blank at 882 nm. The amount of inorganic phosphate was calculated using a standard graph prepared by using  $K_2HPO_4$ .

#### *Estimation of Organic phosphate*

Organic phosphate was estimated by subtracting inorganic phosphate from the total phosphorus.

#### *Estimation of Hardness*

To 10ml of filtered sample 0.2ml of buffer solution and 50mg of Erichrome black T were added. The solution turned wine red. This was treated against EDTA (0.01 M) solution. At the end point, colour change from winered to blue was noted.

#### *Estimation of Calcium*

To 10 ml of filtered sample, 0.4 ml of 0.1 N NaOH and 50 mg of murexide indicator were added. This was titrated against EDTA (0.01 M) solution until the pink colour solution changed to purple.

#### *Estimation of Magnesium*

Amount of magnesium was estimated by the following methods

$$\frac{Mg^{++} (mg l^{-1})}{\text{Volume of sample}} = \frac{Y - X \times 400.8}{\text{EDTA used in calcium determination for the same volume of the sample}}$$

Where, Y=EDTA used in hardness determination, X = EDTA used in calcium determination for the same volume of the sample.

#### *Estimation of chloride*

To 10ml of filtered sample in an Erlenmeyer flask, few drops  $K_2Cr_2O_4$  were added and titrated against silver nitrate (0.0141 N) solution until a persistent red

tinge colour appeared. The amount of chloride was calculated as follows.

#### *Determination of Biological Oxygen Demand*

pH of the water sample was adjusted to neutrality using 1 N acid or 1 N alkali. BOD bottles were filled with water sample without air bubbles. One ml of allylthiourea was added to each bottle. Dissolved oxygen content for 3 of the bottles were estimated by Winkler's method (DO estimation) and the mean of the three bottles were taken (D1). Rest of the bottles (3) was incubated at 27°C in a BOD incubator for 3 days. DO concentrations were estimated after 3 days in these incubated bottles and the mean was calculated (D2).

#### *Chemical Oxygen Demand (COD)*

10ml of sample in 100ml conical flask was taken in triplicate. A blank was run for each using COD free water (Sample is diluted when COD is more than 16  $mg l^{-1}$ ). The flasks were kept in boiling water bath for 1 hour after adding 1ml of 0.1 N  $KMnO_4$  solutions. After cooling for 10 minutes, to each flask, 1ml of KI solution and 2ml of  $H_2SO_4$  were added. Then the sample was titrated with 0.1N sodium thiosulphate and 1ml of starch was added when the solution becomes pale yellow. Titration was continued until the colour disappeared completely. Finally the titre values were noted.

#### **Results and Discussion**

For the present study effluent sample was collected from dye industry. From the sample fungal species were isolated and recorded.

#### *Fungal flora*

Totally 13 species of fungi belong to 8 genus from the effluent were recorded (Table 1). Among the genus *Aspergillus* was recorded as dominant genus with 5 species such as *A.niger*, *A.flavus*, *A.nidulans*, *A.luchensis* and *A.terreus*. The genus, *Trichoderma* was recorded two species (Table 1). The *Trichoderma* such as *Trichoderma sp* and *T. viride* were recorded. Among remaining genus such as *Penicillium*, *Helminthosporium*, *Cladosporium*, *Fusarium*, *Geotrichum* and *Rhizopus* were recorded single species each (Table 1).

#### *Physico-chemical characteristics of effluent pH*

The pH of the effluent was acidic in nature. It was recorded 7.2 initially and the fluctuation of pH was recorded during experimental period (Table 2).

#### *Free CO<sub>2</sub>*

The free  $CO_2$  level was 32  $mg l^{-1}$  initially (Table 2). It was brought down from 15<sup>th</sup> day onwards in both free and immobilized *Aspergillus* treated



#### effluent Alkalinity

The carbonate alkalinity was completely absent in the effluent. Whereas bicarbonate alkalinity was recorded. Initial bicarbonate level in the effluent was  $108 \text{ mg l}^{-1}$  (Table 2). It was reduced to nearly 80% in both free and immobilized *Aspergillus* treated effluent when compared to control.

#### Dissolved Oxygen

After treatment of effluent on 15<sup>th</sup> day the DO level was raised to  $2.2 \text{ mg l}^{-1}$  in and  $1.72 \text{ mg l}^{-1}$  (Table 2) in free and immobilized *Aspergillus* treated effluent respectively.

#### Nitrate, Nitrite and Ammonia

The level of Nitrate, Nitrite and Ammonia were recorded  $90$ ,  $48$  and  $41 \text{ mg l}^{-1}$  initially (Table 2). The all the nitrogen was reduced nearly 50 in both free and immobilized *Aspergillus* treated effluent on 15<sup>th</sup> day (Table 2).

#### Total, inorganic and Organic phosphates

The levels of all forms phosphates were recorded initially in the effluent. Among the phosphates, the total phosphate level was recorded as maximum than the inorganic and organic phosphate. It was reduced on 15<sup>th</sup> day in both free and immobilized *Aspergillus* treated effluent (Table 2).

#### Calcium

The level of calcium was recorded  $83 \text{ mg l}^{-1}$  initially (Table 2). It was brought down from 5<sup>th</sup> day onwards. On 15<sup>th</sup> day it was recorded nearly 45% removal in immobilized *Aspergillus* treated effluent.

#### Magnesium

The level of magnesium hardness was recorded initially  $56 \text{ mg l}^{-1}$ . There was similar reduction as in calcium level on 15<sup>th</sup> day in both free and immobilized *Aspergillus* treated effluent (Table 2).

#### Chloride

Chloride level was recorded initially as  $49 \text{ mg l}^{-1}$  (Table 2). On 15<sup>th</sup> day the calcium was  $44 \text{ mg l}^{-1}$  in immobilized *Aspergillus* treated effluent. Similarly  $40 \text{ mg l}^{-1}$  recorded in free cells of *Aspergillus* treated effluent.

#### BOD and COD

The level of BOD was recorded maximum as  $340 \text{ mg l}^{-1}$  initially (Table 2). It was reduced nearly 65%

in the effluent treated with *Aspergillus* when compared to control. The high amount of COD was recorded initially ( $620 \text{ mg l}^{-1}$ ) (Table 2).

It was reduced in their level in *Aspergillus* treated effluent. In general all the form pollutants removed in *Aspergillus* treated effluent. When compared to control. Similarly immobilized *Aspergillus* treated effluent showed more removal than free cells of *Aspergillus*.

The abundant variation in the types of all living organisms taken together in any geophysical area is called biodiversity. With reference to large organisms (plant, animals etc.) the distinctive morphological and anatomical features visible to the naked eye enable us to identify the different genera and species and, therefore; we can easily assess the extent of their diversity. However, most fungi are not visible to the naked eye and they need to be studied under the microscope. The culturable fungi can be identified provided they sporulate but the biotrophs which are not amenable to culturing in growth media go unnoticed. It is, therefore, imperative that there is a huge gap in our knowledge of the diversity of all microbes including fungi. Fungi are better known than bacteria, viruses and other smaller forms of life such as viroids. There are about 72,000 named species and new species are being added at the rate of about 1500 each year. The total number of fungal species both known and unknown are estimated at 250,000 (Hawksworth & Kirsop, 1988; Heywood, 1995). The 72,000 known species exist in different culture collection centres in the form of several subspecies, pathovars and strains, the total cultures going up to 170,000 (Bennett & Faison, 1997).

The mycoflora of unique ecological niches have some common features and it will be very rewarding to explore fungal diversity in habitats such as thermophilic environments, e.g., hot springs, thermal vents, sun-heated soils, compost pits, self-heated coal refuse piles, steam line discharge sites, etc. The thermophilic fungi and bacteria grow at temperatures between  $40\text{--}60^\circ\text{C}$  are the sources of thermostable enzymes. Other ecological niches to be explored are western ghats, the Himalayan ranges, marine ecosystems, mangroves, coral reefs, sand dunes, industrial effluent contaminated soils, refineries, activated sludge, insects and several other natural sources. Endophytes of plants and animals are very meagerly understood.

Hence, the present study was undertaken to know the fungal species in industrial effluent. For the present investigation effluent sample was collected from leather industry. The fungal species were isolated and identified from the effluent sample were recorded (Table.1).

Totally 13 species of fungi belongs to 8 genus from



the effluent (Table 1). Among the genus *Aspergillus* was recorded as dominant genus with 5 species such as, *A. niger*, *A. flavus*, *A. nidulans*, *A. luchensis* and *A. terreus*. Similar work was done by Carroll and Wicklow (1992). They isolated 17 species of fungi in the household wastewater in which *Aspergillus* was the dominant group of fungi.

In this present study two species of *Trichoderma* such as *Trichoderma* sp and *T. viride*. The present results were in conformity with the earlier reports of Watts *et al.*, (1999). They isolated and recorded three species of *Trichoderma* from automobile industrial effluents. In this present study *Penicillium*, *Helminthosporium*, *Cladosporium*, *Fusarium*, *Geotrichum* and *Rhizopus* were recorded single species each. These types of fungal species always occur in industrial effluent (Bennett & Faison, 1997).

The ability of fungi to transform a wide variety of hazardous chemicals has aroused interest in using them in bioremediation (Alexander, 1994). The white rot fungi are unique among eukaryotes for having evolved nonspecific methods for the degradation of lignin; curiously they do not use lignin as a carbon source for their growth (Kirk *et al.*, 1976). Lignin degradation is, therefore, essentially a secondary metabolic process, not required for the main growth process. Lamar *et al.*, (1993) compared the abilities of three lignin-degrading fungi, *Phanerochaete chrysosporium*, *P. sordida* and *Trametes hirsuta* to degrade PCP (Pentachlorophenyl) and creosote in soil. Inoculation of soil with 10% (wt/wt) *Phanerochaete sordida* resulted in the greatest decrease of PCP and creosote. *P. sordida* was also most useful in the degradation of PAHs (Polycyclic aromatic hydrocarbons) from soil. Davis *et al.*, (1993) showed that *P. sordida* was capable of degrading efficiently the three ring PAHs, but less efficiently the four-ring PAHs.

The present study was carried out to treat the leather industry effluent by using the dominant species of *Aspergillus*. In this study all the pollutants were reduced in their level in both unsterilized and sterilized effluent due to the inoculation of *Aspergillus*. The similar work was done by Thanh and Simard (1973). They reported treatment of wastewater by fungal species. While maximizing reduction of phosphate, ammonia, and organic matter. The studies were conducted in batch 500-ml baffled culture shake flasks containing 150 ml of sterile wastewater. Reported phosphate

removal ranged from 12 to 100%, total nitrogen removal from 22 to 93%, ammonia nitrogen from 27 to 90%, and COD removal from 0 to 72%.

Hiremath *et al.*, (1985) performed a similar study except they tested seven fungal species isolated from a wastewater stabilization pond. The major goal of the study was to maximize biomass production of fungi as a food source for animal or human consumption. The trials were conducted in 2-L conical flasks containing

1.5 L of sterile fresh wastewater. Flasks were inoculated with pure cultures of fungi and incubated at room temperature for 10 days. The culture flasks were gently agitated twice a day. The study reported BOD removal between 53 and 72%, phosphate removal from 34 to 77%, and ammonia nitrogen removal between 49 and 77%. Due to the experimental design, the cultures were most likely completely anaerobic for the entire 10-days incubation.

Both studies were performed under non-ideal conditions with process optimization of the parameters reported. Although the data showed fungi will treat wastewater, it does not give any indication of the maximum removal efficiencies. However, both studies indicated promising results for removal of nitrogen and phosphorus. Additional evidence, although not conclusive, that fungi are capable of wastewater treatment was found during the study of onsite sphagnum peat wastewater treatment system. Brooks (1988) found excellent removal of BOD, organic nitrogen, and ammonia nitrogen (90 to 95% and 95 to 99%, respectively). Based on standard bacterial and fungi enumeration techniques, the ratio of fungi to bacteria was 8:1 during winter and 4:1 during the summer. This led to hypothesize that fungi had a large role in removal of nitrogen from the wastewater. However, due to the use of standard enumeration techniques it is very likely the populations were underestimated due to the fastidious nature of environmental microbes. Furthermore, a direct comparison ratio of fungi to bacterial colony forming units is not appropriate. This ratio employs the flawed assumption that each bacteria and fungi is equal in degradation rate and range (BOD, ammonia nitrogen, and organic nitrogen) of the wastewater. A different experimental design and enumeration is needed before comparisons can be made.

Whenever bioremediation figures as the topic of discussion, bacterial agents come into focus and fungi are much less studied. One should realize, however, the greater potential of fungi by virtue of



their aggressive growth, greater biomass production and extensive hyphal reach in soil. More research will be focused in future on using the diverse fungal flora for bioremediation. The work

on the fungal diversity in the chemical industrial effluents and promising fungi *Aspergillus flavus* that can degrade various pollutants in these effluents.

- hydrocarbons" by *Pleurotus ostreatus* in soil. *Applied and Environmental Microbiology*, 2000: 66, 6, 2471-2478.
7. Balice V, Boari G, Cera O and Abbaticchio P. "Indagine analitica sulle acque di vegetazione. Inquinamento", 1982: 7, 49-53.
  8. Belinky P A, Flinshtein N, Lechenko S, Gepstein S, and Dosoretz C. "Reactive oxygen species and induction of lignin peroxidase in *Phanerochaete chrysosporium*". *Applied and Environmental Microbiology*, 2003: 69, 11, 6500-6506. Castro, A.I.R.P, Eytuguin, D.V and Xavier, A.M.R.B." Degradation of biphenyl lignin model compounds by laccase of *Trametes versicolor* in the presence of 1- hydroxybenzotriazole and heteropolyanion". [SiW11VO40]<sup>5-</sup>. *Journal of Molecular Catalysis B: Enzymatic*, 2003: 22, 13-20.
  9. YN Jolly, A Islam, AlMustafa. "Characterization of dye Industry Effluent and Assessment of Its Suitability for Irrigation Purpose". *Journal of Bangladesh Academy of Sciences*, 2009: 33, 1

**Table 1: Physicochemical characteristics of effluent**

| S.No | Parameters           | Initial | 5 <sup>th</sup> Day |         |      | 10 <sup>th</sup> Day |         |      | 15 <sup>th</sup> Day |         |      |
|------|----------------------|---------|---------------------|---------|------|----------------------|---------|------|----------------------|---------|------|
|      |                      |         | Control             | Treated |      | Control              | Treated |      | Control              | Treated |      |
|      |                      |         |                     | FC      | IMC  |                      | FC      | IMC  |                      | FC      | IMC  |
| 1.   | pH                   | 7.2     | 6.8                 | 6.8     | 6.7  | 6.8                  | 6.6     | 6.7  | 6.8                  | 6.6     | 6.7  |
| 2.   | Free CO <sub>2</sub> | 32      | 30                  | 28      | 24   | 30                   | 22      | 20   | 28                   | 12      | 10   |
| 3.   | Alkalinity           | 108     | 106                 | 78      | 96   | 106                  | 62      | 48   | 104                  | 20      | 21   |
| 4.   | Dissolved Oxygen     | 1.35    | 1.72                | 1.74    | 1.37 | 1.72                 | 2.2     | 1.37 | 2.2                  | 2.2     | 1.72 |
| 5.   | Nitrate              | 90      | 90                  | 82      | 70   | 90                   | 68      | 59   | 88                   | 42      | 52   |
| 6.   | Nitrite              | 48      | 48                  | 46      | 40   | 46                   | 36      | 32   | 44                   | 26      | 28   |
| 7.   | Ammonia              | 41      | 41                  | 32      | 30   | 40                   | 26      | 22   | 39                   | 16      | 16   |
| 8.   | Total Phosphorus     | 147     | 146                 | 136     | 130  | 144                  | 126     | 168  | 142                  | 80      | 78   |
| 9.   | Inorganic            | 83      | 82                  | 78      | 23   | 80                   | 72      | 62   | 78                   | 42      | 42   |
| 10.  | Organic              | 64      | 64                  | 58      | 57   | 62                   | 54      | 44   | 60                   | 38      | 30   |
| 11.  | Calcium              | 83      | 82                  | 76      | 68   | 80                   | 70      | 58   | 78                   | 50      | 49   |
| 12.  | Magnesium            | 56      | 55                  | 50      | 50   | 54                   | 42      | 45   | 54                   | 22      | 37   |
| 13.  | Chloride             | 49      | 44                  | 48      | 44   | 49                   | 46      | 42   | 48                   | 44      | 40   |
| 14.  | BOD                  | 340     | 320                 | 250     | 260  | 310                  | 220     | 180  | 310                  | 160     | 110  |
| 15.  | COD                  | 620     | 610                 | 520     | 500  | 600                  | 420     | 380  | 580                  | 300     | 220  |